

SYNOPTIC ANALYSES, 5-, 2-,  
AND 0.4-MILLIBAR SURFACES  
FOR JANUARY 1972  
THROUGH JUNE 1973



CASE FILE  
COPY



NATIONAL AERONAUTICS AND SPACE ADMINISTRATION  
and  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION



SYNOPTIC ANALYSES, 5-, 2-,  
AND 0.4-MILLIBAR SURFACES  
FOR JANUARY 1972  
THROUGH JUNE 1973

A joint NASA-NOAA publication prepared for NASA Wallops Flight Center  
by the Staff, Upper Air Branch, National Weather Service, Camp Springs, Maryland  
under Wallops Flight Center Purchase Order Number P-55946(G)



*Scientific and Technical Information Office*

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

*Washington, D.C.*

1975

## CONTENTS

INTRODUCTION . . . . .	1
PROCESSING OF ROCKETSONDE DATA . . . . .	2
SATELLITE DATA . . . . .	6
ANALYSIS PROCEDURE . . . . .	6
DISCUSSION OF THE JANUARY 1972 - JUNE 1973 CIRCULATION . . . . .	8
ACKNOWLEDGMENTS . . . . .	11
REFERENCES . . . . .	12
WEEKLY 5-, 2-, and 0.4-MB SYNOPTIC CHARTS . . . . .	29

## LIST OF ILLUSTRATIONS

### Figure

1. Temperature measured at Ascension I., Ft. Sherman and Thumba . . . . .	14
2. Relationships between radiances and thickness and between radiance and temperature . . . . .	15
3. Analyzed values from weekly charts at Fort Greely . . . . .	16, 17
4. Analyzed values from weekly charts at Fort Churchill . . . . .	18, 19
5. Analyzed values from weekly charts at Volgograd . . . . .	20, 21
6. Analyzed values from weekly charts at Wallops Island . . . . .	22, 23
7. Analyzed values from weekly charts at White Sands . . . . .	24, 25
8. Analysed values from weekly charts at Antigua . . . . .	26, 27
9. Station model and reporting rocket stations . . . . .	28

# SYNOPTIC ANALYSES, 5-, 2-, AND 0.4-MILLIBAR SURFACES

FOR JANUARY 1972 THROUGH JUNE 1973

By Staff,<sup>1</sup> Upper Air Branch  
NOAA, National Weather Service, National Meteorological Center

## SUMMARY

Data from meteorological rocketsonde and satellite radiance measurements have been employed to analyze a series of high-altitude synoptic charts. The methods employed for processing the various types of data and the analysis procedure used are described.

Broad-scale analyses for the Northern Hemisphere 5-, 2-, and 0.4-mb surfaces are presented weekly from September through April and monthly from May through August during the period January 1972 through June 1973. A brief discussion of the height and temperature fields is also given. Circulation and temperature changes associated with a minor stratospheric warming in January and February 1972 and a major stratospheric warming in January and February 1973 are among the discussion items.

## INTRODUCTION

This report is the sixth in a series of constant-pressure charts for the upper stratosphere and lower mesosphere. Charts for 1964-68 (refs. 1, 2, 3, 4, 5) were analyzed at weekly intervals (ref. 6) and were based on meteorological rocketsonde and very high level rawinsonde data obtained over Northern America and adjacent ocean areas (ref. 7).

Since 1972 (pursuant to a bilateral agreement between the United States and the Soviet Union), data from meteorological rocketsonde stations of the Western and Eastern Meridional Networks have been exchanged on a regular basis (ref. 8). Recently available data from Heiss Island and Volgograd (Soviet Union) and Thumba (India) together with other data from Ryori (Japan) and El Arenosillo (Spain) have made it possible to extend the analyses over Eurasia. Figure 9 shows the locations of the 20 meteorological rocket launch sites over the Northern Hemisphere for which data were available. Because of improvements in the methods of routine transmission and receipt of rocketsonde data by the teletypewriter-coded ROCOB messages, charts could be constructed from these data rather than delaying until publication of the rocketsonde observations.

---

<sup>1</sup>Personnel actively engaged in this report were Manager, F. Finger; Coordinator and Analyst, M. E. Gelman; Analysts, R. Nagatani and J. Laver; Research Consultants, R. S. Quiroz and A. J. Miller; and data processing, D. Griffith.



Recently, data from satellite vertical temperature sounding instruments also have become available (see section on satellite data, below). These remote observations, when used together with the expanded number of in situ observations, provide for greatly improved data coverage over the entire hemisphere.

The present series of analyses at the 5-, 2-, and 0.4-mb surfaces (approx 36, 42, and 55 km, respectively) represent the broad-scale synoptic conditions over the Northern Hemisphere on each Wednesday during January 1972 through June 1973. Charts are presented weekly during the winter months (October through March) and transition periods (April and September) and monthly during summer (May through August). The monthly frequency appears sufficient to depict the slow, broad-scale changes that occur in summer. Small-scale time and space changes and planetary waves of small amplitude (ref. 9) are observed during summer, but the horizontal resolution of the analyses in most areas would not be sufficient to depict them. During winter and the transition periods, however, larger changes are evident from one week to the next. Sometimes, large changes occur within a day or two, and these may be inferred from the sequence of up to three observations plotted at each rocket station. Thus the user may expand the utility of the weekly charts by noting any large changes during the week in plotted temperature or wind direction. These changes frequently indicate movements of the synoptic systems depicted in the Wednesday charts.

Despite the omission of small-scale details, the maps are very useful for a number of applications such as determining the trajectory of constant-level balloons, relating variations in infrasound propagation to circulation changes, and providing a data base for evaluating environmental effects on aerospace vehicles. In addition, users have pointed to the increasing utility of these maps in studies of stratospheric-ionospheric interaction, for verification of the performance of numerical circulation models, and in various other research efforts.

#### PROCESSING OF ROCKETSONDE DATA

Temperature, height and wind information derived from routine meteorological rocketsonde observations comprised the basic data for analyses at the 5-, 2-, and 0.4-mb levels. Rocketsonde information used for this project were obtained from teletype coded ROCOB messages (WMO code FM39.E ROCOB and FM40.E ROCOB ship). These data were usually transmitted from each rocketsonde station within one day of the observation. Data from Heiss Island, Volgograd and Thumba (The three stations in the Northern Hemisphere associated with the Eastern Meridional Network), were available in ROCOB format within one week of observation time.

The ROCOB message provides data in the form of temperature and wind versus geometric height. The vertical coordinate must be transformed from height to pressure so that temperature, height, and wind information may be extracted at the desired levels.

The procedure for calculating pressure and extracting the required information for the 5-, 2-, and 0.4-mb levels was as follows:

- a. Pressure was calculated at each ROCOB reported level by a computer program which integrated the hydrostatic equation starting at a base level

near 50 mB. Temperature and height data obtained from a nearby rawinsonde station were used as the reference-level data. The geopotential heights and temperatures at the 5-, 2-, and 0.4-mb levels were interpolated in this computer program.

b. Wind direction and wind speed (in knots) were interpolated manually at the calculated height of each analysis level. When temperature data were not available for a particular sounding, the wind information was extracted at individually estimated heights of the 5-, 2-, and 0.4-mb levels.

c. Time-height diagrams were plotted for each of the 20 rocketsonde stations. The temperature and wind information on these diagrams provided valuable verification of the sequence of meteorological changes. Erroneous or questionable data could be quickly isolated in this manner.

d. From the wind information plotted on the time-height diagrams, thermal winds were determined for approximately 6-km layers surrounding each analysis level. A somewhat subjective approach was necessary because the calculations were performed manually and there were sometimes rapid wind oscillations with height. Usually an unambiguous direction for the thermal wind could be determined.

e. Special adjustments were made to the 0.4-mb data because ROCOB data normally contain uncorrected temperatures. Thermistor rocket soundings generally require correction above about 40 km. Temperature adjustment values are given in Table 1a, b, and the basis for these adjustments is described below.

(i) For the U.S. Datasonde instrument used at the majority of Northern Hemisphere stations, corrections have been derived by theoretical and laboratory methods (ref. 10). The corrections account for aerodynamic heating (which depends mainly on the fall velocity of the sensor), thermal lag, emitted and absorbed radiation, and electrical heating. Precise corrections based on these factors vary from sounding to sounding and can only be calculated from auxiliary rocketsonde information. For our purposes, it was deemed satisfactory to derive a set of corrections based on average sensor fall velocities and for sunlight and darkness conditions (Table 1c).

(ii) Application of the average corrections at 0.4 mb is not sufficient to achieve compatibility between U.S. and U.S.S.R. ROCOB temperatures. Indirect comparisons between U.S.S.R. and U.S. uncorrected rocketsonde temperatures indicate large differences increasing with height above 45 km. An example of such an indirect comparison is shown in Figure 1 for the U.S. rocketsonde station at Fort Sherman, C.Z., and Thumba, where the U.S.S.R. rocketsonde is used. Although these stations are geographically distant, they are within one degree of the same latitude circle, and there is no evidence (satellite data, thermal winds etc.) supporting the existence of a real standing wave in temperature throughout the year of the magnitude inferred from the data from these stations. Comparisons were also made for the U.S. station at Thule and the U.S.S.R. station at Heiss Island, both at latitudes near 80°N, with results strikingly consistent with those in Figure 1.

TABLE I. - THEORETICAL CORRECTIONS AND EMPIRICAL TEMPERATURE ADJUSTMENTS (°C) DERIVED TO OBTAIN COMPATIBILITY BETWEEN U.S. AND U.S.S.R. ROCKETSONDE MEASUREMENTS

KM	(a)		(b)		(c)		(d)		(e)	
	U.S. Adjustment Day	U.S. Adjustment Night	U.S.S.R. Adjustment	Average Theoretical U.S. Correction	Day	Night	U.S.S.R. Minus U.S. Empirical	U.S. Difference Night Minus Day		
58	-16	-4	9	-6.0	-3.6		-25	-12		
55	-8	-2	7	-4.2	-2.4		-15	-6		
50	-5	-1	2	-2.5	-1.4		-7	-4		



Although differences between the U.S. and U.S.S.R. temperature data (shown in Table 1d) were derived indirectly, there appears to be little doubt that temperatures from the U.S.S.R. observations are significantly lower than those reported by the U.S. instrument. Correction of the U.S. reports by values shown in Table 1c would reduce the incompatibility only partially.

Meanwhile, it is known that very large corrections are applied by the Soviet scientists in order to compensate for very large fall velocities. Thus the residual uncertainty in their corrected temperatures near 55 km may possibly exceed the uncertainty in corrected U.S. soundings.

(iii) There is a complicating factor when trying to rationalize the seeming incompatibility between U.S. and U.S.S.R. temperature data. This factor is the temperature change brought on by the real diurnal temperature variation of the atmosphere. If all rocket observations were taken at the same local time, this would not pose any problems. However, The U.S.S.R. rockets are launched primarily during nighttime hours while U.S. rockets normally are launched near noon. Thus, the amplitude and phase of the diurnal temperature oscillation are undoubtedly an important factor. In order that a consistency in time be maintained in the analyses, adjustments must include this factor. The adjustments for U.S. instruments are based on an experiment held at Wallops Island, Virginia in October and November 1972. A series of mid afternoon, pre-sunset and post-sunset observations (uncorrected) indicate the day-night differences shown in Table 1e. Real diurnal temperature change as well as instrumental radiation error are the principal factors for these differences. Thus a total adjustment to the U.S. measurements taken during the day to reduce them to ambient nighttime values is the sum of the empirical day-night difference and the theoretical nighttime correction. However, even this total adjustment (Table 1a) to the U.S. measurements does not account for the entire observed U.S.—U.S.S.R. temperature differences. The remaining portion has been assigned as an adjustment to the U.S.S.R. data (Table 1b). All adjustments to U.S. reports result in temperatures lower than reported, while U.S.S.R. adjustments result in higher than reported temperatures. The calculated 0.4-mb heights from U.S. instruments were lowered by 200 gpm. This represents the approximate integrated effect of the temperature measurement being too high from 0C degrees at 2 mb to 8C degrees at 0.4 mb. The 0.4-mb heights from U.S.S.R. instruments were raised by approximately 100 gpm.

The rocketsonde data, temperature (C°), height (geopotential meters), wind direction and speed (knots), were plotted on a polar stereographic map base. On the charts presented for publication three observations closest to Wednesday are shown when available for each station. Reported heights and calculated thermal winds have been omitted for the sake of legibility. The station model chart (Fig. 9) illustrates the symbols used to distinguish data obtained on Wednesdays from off-time data.

## SATELLITE DATA

For January to April 1972, Nimbus 4 Satellite Infrared Spectrometer (SIRS) and Selective Chopper Radiometer (SCR) (ref. 11) data were used for the analyses at 5, 2, and 0.4 mb. Nimbus 5 SCR (ref. 12) and NOAA 2 Vertical Temperature Profile Radiometer (VTPR) (ref. 13) data were used for the 1973 analyses. The method of using the remotely sensed temperature information for determining stratospheric thickness was developed by Quiroz and Gelman (ref. 14). In brief, the radiant energy sensed by a satellite instrument in any spectral band is representative of the weighted temperature from a substantial layer in the atmosphere. Single channel relationships were derived relating satellite measured radiances and the radiosonde-rocketsonde thickness (or mean temperature) between the 100- to 5-mb, 100- to 2-mb, and 10- to 0.4-mb levels. These thickness relationships are shown in Figure 2a. A table was then constructed relating observed radiance with thickness values at 320 meter intervals for each of these atmospheric layers. Using this table, the radiance map containing 24 hours of satellite data was analyzed for the appropriate radiance interval, with the radiance lines then being converted to thickness values. This thickness field was then graphically added to the height field given by the objectively-analyzed chart produced at the National Meteorological Center (ref. 15) for the lower level (100 mb for the 5- and 2-mb charts, and 10 mb for the 0.4-mb charts). The resultant field was used as a first guess for the 5-, 2-, and 0.4-mb height chart.

Relationships between radiance and temperature at 5, 2, and 0.4 mb were also sought. A weaker physical relationship exists between radiance and the temperature at any particular level than exists between radiance and the mean layer temperature or thickness. However, it was found that SCR channel B and SCR channel A specified the temperature at 5 and 2 mb to a good approximation (RMS error, approximately 8°C). These relationships are shown in Figure 2b. The temperature patterns obtained by relabeling the appropriate radiance charts were then used as a first guess in deriving the 5- and 2-mb temperature analyses.

## ANALYSIS PROCEDURE

The analysis procedure consisted of obtaining first guess temperature and height fields and then adjusting these fields to conform with the rocketsonde wind, height, temperature and thermal wind information. When satellite data were available, first guess fields were obtained using the methods discussed in the previous section. If satellite data were not available, conventional techniques were used to construct the temperature field, first at 5 mb; then differential analysis methods were used to obtain the height fields. Once the 5-mb fields were completed, the fields at 2 mb, then at 0.4 mb were built in a similar manner.

The analysis system consisted of the following steps:

- a. Isotherms were derived using as primary information the plotted rocketsonde temperatures. Data acquired for the entire week were examined to determine the synoptic changes that took place during the week. Thus, conditions pertaining to Wednesday, the analysis day, were deduced. The SCR radiance data, when available, were especially useful in providing a first approximation to the temperature fields. Computed thermal winds were

very useful in determining horizontal temperature gradients and the relative location of warm and cold areas. Time-height sections of temperature were consulted as a further aid in deriving the isotherms.

b. When satellite data were not available, a first guess height field was derived by differential analysis. A mean temperature field representative of the layer between the previously analyzed lower surface and the selected surface was derived graphically. This mean field represents a geopotential thickness which, when added to the lower level height field, yields a smooth, conservative first approximation to the contour pattern at the upper surface.

c. Reported winds and computed heights for individual stations were then used to adjust the first approximation of the contour field, assuming geostrophic flow. Winds were accorded the highest priority for this adjustment. When large adjustments were made to the contour field, the temperature field was then adjusted to maintain hydrostatic consistency.

d. The analyses were reviewed for vertical and temporal consistency. For example, circulation centers, ridges, and troughs were examined with the aid of all available data to verify vertical slope and movement with time. Time-height sections and height-change charts were especially useful for those purposes.

The above procedures appear to produce excellent results at 5 mb and 2 mb, and were successfully applied to obtain the 0.4-mb charts. Generally only slight adjustments of the first-approximation height fields were necessary at the 5- and 2-mb levels. However, rather formidable analysis problems were evident at the 0.4-mb level.

One problem arises from the apparent intersection of the stratopause with the 0.4-mb level. Because the normal stratospheric temperature inversion ceases at the stratopause level, the graphical method for obtaining mean temperature, which depends on the existence of a linear profile, is no longer valid. Large adjustments must be made, especially at lower latitudes, in the graphically derived height field to conform with the computed height at each station. This difficulty is largely overcome when satellite data are available for use in deriving first guess fields.

Another difficulty was the apparent occurrence of large day-to-day temperature changes, at times exceeding  $10^{\circ}\text{C}$  (ref. 16), and persistent oscillations in many wind profiles. In most cases, deviations of reported temperatures and winds from one another could be accounted for by identifiable rapid large-scale synoptic changes. Sometimes rocketsonde reports within a few hours of each other at a single station exhibited temperature changes of  $5^{\circ}$  to  $7^{\circ}\text{C}$  over a limited height interval near the stratopause. It has been shown that such small-scale subsynoptic changes are possible at these levels (ref. 17). Thus, some intermediate value was chosen for analysis to represent the value on the analysis day. Occasionally, it was impossible to make a reasonable reconciliation of reported station values.

Although careful consideration of high-level data allows a broad-scale depiction of circulation patterns up to 0.4 mb, the sparsity of reports requires increasing subjectivity as the analysis proceeds to this level. The justification for some analyses depends on the interpretation of the limited amount of data in such a way as to portray a coherent sequence of synoptic events. In spite of these factors, surprisingly



little alteration in the principal features of the circulation and temperature distribution shown in the final analysis can be made without inordinately violating some of the data. In general the contours and isotherms depicted are felt to be good approximations to the flow and temperature patterns at this level. Even so, the same degree of accuracy that is found customarily in the analysis of charts at lower levels should not be expected.

A contour interval of 320 geopotential meters was used throughout the year. In addition, intermediate dashed contours were used to outline areas of relatively weak gradient, especially during the spring and fall changeover periods. Isotherms were drawn and labeled at 5°C intervals.

## DISCUSSION OF THE JANUARY 1972 - JUNE 1973 CIRCULATION

The temperature and circulation patterns during January and February 1972 are of considerable interest because of the large scale changes associated with warmings in the upper stratosphere. However, these stratospheric warming events were not major warmings with respect to their influence on the circulation of the lower stratosphere.

In the first week of January, a significant warm area was apparent over much of the Northern Hemisphere, extending from eastern North America at mid-latitudes and spiralling around Eurasia northeastward to eastern Siberia. The arctic cold center became elongated at 5 mb and developed into a double center at 2 mb, while warm air prevailed over the polar area at 0.4 mb. Strong temperature gradients characterized the situation at each level; as much thermal contrast was evident longitudinally as there was with latitude. The height field also reflected a strongly disturbed wintertime situation, with an anticyclonic circulation in the northern Pacific area at 5 mb intensifying and sloping westward at higher levels. The Atlantic anticyclonic circulation remained somewhat less intense than the Pacific feature. The dominant polar vortex was constricted at 5 mb and became elongated between the two anticyclonic circulations at higher levels. Maximum westerly winds approaching 300 knots were reported by rocketsonde stations located in the area associated with the extreme pressure contrasts. The large wind shears associated with these strong winds are further evidence of the intense thermal gradients and the movements and slope with height of the warm-air systems.

During the second week in January a warm area at 2 mb was evident over the polar region in a constricted channel extending from southern Asia and over the Pole to Hudson Bay. This warm area was a persisting feature throughout the next two months at the 2 and 0.4 mb levels, with two intense cold areas located over Europe and the northern Pacific region. The next three weeks remained very strongly disturbed at all levels with the two opposing anticyclones over the Pacific and the Atlantic oscillating both in intensity and position. At the end of January, a region of warm air appeared west of Japan at 5 mb.

On February 9 at 5 mb, the Atlantic anticyclone showed a significant increase in intensity, and the polar vortex became strongly elongated between Atlantic and Pacific anticyclones. The region of warm air which first appeared west of Japan at the end of January made a rapid incursion into the polar region by February 16, accompanied by a marked warming at

5 and 2 mb. The temperature in the warmest areas at 5 mb increased from -20C the previous week to +10C, with temperatures near the Pole increasing about 50C degrees. At 2 mb, the warmest areas increased from -5C to +20C, with polar temperatures increasing about 25C degrees. The changes at 0.4 mb were much less pronounced.

The sudden warm air incursion was followed on February 23 by a correspondingly sudden decrease in intensity of the warm air in the middle stratosphere as the anomaly affected lower levels. The Pacific anticyclone moved to northern Alaska in partial response to the warming at lower levels. On March 1, the polar vortex at 5 mb, split into two cells with the warmest air cooling still further. By March 8, westerly circulation was once again dominant over the northern latitudes, with weaker anticyclonic circulation still evident at middle and low latitudes. Throughout all this activity, however, there was no major reversal of circulation patterns in the lower stratosphere; hence the label of "minor warming" for this period of activity. The temperature gradients became less intense in March as the springtime changeover proceeded. By mid-April, at the 5-mb level, anticyclonic circulation and warm air dominated the polar region, with the reversal in temperature gradient and circulation occurring at higher levels within two weeks. The westerly winds associated with cyclonic circulation were displaced equatorward, with a cellular structure quite evident during this period.

As the summer months approached, the easterly circulation became the dominant feature and the polar anticyclone intensified in response to the solar radiational heating. Maximum temperatures as well as maximum easterly circulation were observed in mid-July. Cooling was evident thereafter in the polar latitudes as perturbations in the easterly circulations developed. A bicellular anticyclonic structure was evident at polar latitudes on the charts for August 16, and by September 5 the developing cyclonic system had formed. As the polar cyclone progressively cooled and intensified, the easterly circulation moved equatorward in the form of cellular ridges. Throughout October, November and December, the polar cyclone deepened, accompanied by increasing westerly winds. Temperatures decreased markedly at 5 and 2 mb during this period but a warm region remained in middle latitudes at 0.4 mb. By December, however, the temperature pattern at the 0.4-mb level was characterized by two cold areas and two warm areas from mid to high latitudes. This pattern persisted for several weeks while the 0.4-mb circulation remained strongly zonal. At low latitudes the subtropical ridge was apparent between 5 degrees and 25 degrees north.

A major stratospheric warming occurred during late January and early February 1973. There were indications of minor pulses as early as December 1972 when the two upper-VTPR channels showed significant radiance increases from December 27 to January 8 with the greatest change beginning in late January (ref. 18). Similarly, the maps which were drawn with the aid of the same radiance data show thermal pulses through the end of January, with a major circulation change occurring at all levels on January 31. Throughout the period leading up to the warming event, anomalously warm air was confined mainly at 2 mb and above in a constricted channel observed over the polar region. Evidence of this is provided by rocketsonde temperatures reported for Heiss Island and Thule. Poker Flat observations on January 16-18 show easterlies only at 0.4 mb while Primrose Lake also showed a report of an easterly wind on January 16. This indicates that the thermal anomaly at that time was strong enough for a weak anticyclonic circulation only at that level. A shift of the Aleutian

anticyclone occurred at 5 mb on January 24, extending easterly circulation into the central U.S. but damping quite rapidly with height.

The period January 24 - 31 was an explosive one. Strong easterlies near the Pole at all three levels on January 31 show that the circulation had already reversed. High temperatures evident upward from 100 mb dominated the maps at 5 and 2 mb. Cooler air at 0.4 mb moving toward the polar regions was overriding the warmer air below. At 5 mb, a ridge of warm air in two cells spiralled from low latitudes starting near Mexico, along the Atlantic into Eurasia, over the Pole and into the north Pacific. A region of cold air remained over the U.S. Similarly, at 2 mb, the ridge of warm air also spiralled in towards the Pole with the ridge at 2 mb lying in more northerly latitudes than the one at 5 mb. The circulation at 0.4 mb consisted of an anticyclone near the Pole with a discontinuous trough made up of three cells of low pressures at mid-latitudes and a ridge of high pressure south of the trough. Cooler air was already moving in towards the Pole at this level.

By mid-February, the polar vortex was re-established at the three analysis levels, with cooler temperatures prevailing and conditions remaining somewhat static for the remainder of the period through mid-April. It is noteworthy that at the 10-mb level, as shown by daily charts (ref. 15), a very weak cyclonic circulation prevailed until early March, when a moderately strong polar vortex was established. Minor thermal pulses were evident at 0.4 mb during March-April, but the heights of the 0.4 mb surface were not significantly altered. By April 18, an anticyclonic circulation appeared over the Atlantic, signifying the start of the changeover period. The anticyclonic circulation had become well established by May 16 at 0.4 mb with indications of a more cellular structure at 5 mb. In June, easterlies were prevalent throughout the layer from 5 to 0.4 mb.

The principal features of the circulation during January 1972 to June 1973 at six representative rocket launch sites may be seen in the time sections (Figs 3-8) of analyzed height and temperature values extracted from the weekly charts. The annual trend of the height and temperature values is most pronounced at the middle- and high-latitude stations (Figs. 3-6), with the largest range of values occurring in the more northerly stations. Superimposed on this annual trend, characterized by maximum values in summer and minimum values in winter, are various perturbations. The largest of these perturbations occurred during January 1973 and January 1972 and were associated with the stratospheric warmings mentioned above. Smaller scale disturbances associated with movements of the Aleutian anticyclone were seen throughout the winter period. At the tropical station of Antigua, West Indies Associated State (W.I.A.S.), the range of values is very small, with only very minor perturbations affecting the station (Fig. 8). There is a suggestion, however, of a double maximum at the time of the equinoxes associated with the semiannual variation in temperature and wind and related to the seasonal migration of the subtropical anticyclone as seen in the charts. At the subtropical station of White Sands, N. Mex. (Fig. 7), a combination of the major annual variation and the semiannual variation is apparent.



## ACKNOWLEDGMENTS

We are grateful to D. Wark, M. Weinreb, and H. Woolf of the National Environmental Satellite Service, NOAA for their assistance in the acquisition of satellite radiance data.

Funds for the chart series were provided by the National Aeronautics and Space Administration under the EXAMETNET program, contract P52774(G).

## REFERENCES

1. Staff, Upper Air Branch, National Meteorological Center, "Weekly Synoptic Analyses 5-, 2-, and 0.4-Mb. Surfaces for 1964," ESSA Technical Report WB 2, Weather Bureau, Environmental Science Services Administration, U.S. Department of Commerce, Silver Spring, Md., Apr. 1967, 176 pp.
2. Staff, Upper Air Branch, National Meteorological Center, "Weekly Synoptic Analyses, 5-, 2-, and 0.4-Mb. Surfaces for 1965," ESSA Technical Report WB 3, Weather Bureau, Environmental Science Services Administration, U.S. Department of Commerce, Silver Spring, Md., Aug. 1967, 173 pp.
3. Staff, Upper Air Branch, National Meteorological Center, "Weekly Synoptic Analyses, 5-, 2-, and 0.4-Millibar Surfaces for 1966 " ESSA Technical Report WB 9, Weather Bureau, Environmental Science Services Administration, U.S. Department of Commerce, Silver Spring, Md., Jan. 1969, 169 pp.
4. Staff, Upper Air Branch, National Meteorological Center, "Weekly Synoptic Analyses, 5-, 2-, and 0.4-Millibar Surfaces for 1967," ESSA Technical Report WB 12, Weather Bureau, Environmental Science Services Administration, U.S. Department of Commerce, Silver Spring, Md., Jan. 1970, 169 pp.
5. Staff, Upper Air Branch, National Meteorological Center, "Weekly Synoptic Analyses, 5-, 2-, and 0.4-Millibar Surfaces for 1968," NOAA Technical Report NWS 14, National Weather Service, National Oceanographic and Atmospheric Administration, U.S. Department of Commerce, Silver Spring, Md., May 1971, 169 pp.
6. Finger, Frederick G., Woolf, Harold M., and Anderson, Calvin E., "Synoptic Analyses of the 5-, 2-, and 0.4-Millibar Surfaces for the IQSY Period," Monthly Weather Review, Vol. 94, No. 11, Nov. 1966, pp. 651-661.
7. World Data Center A, Meteorology, Data Report, Meteorological Rocket Network Firings 1964-1968, Vol. I-V, National Weather Records Center, Environmental Science Services Administration, U.S. Department of Commerce, Asheville, N.C., 1967-1971.
8. FXAMETNET, Experimental InterAmerican Meteorological Rocket Network: 8th Annual Meeting Report, Oct. 30-Nov. 3, 1972, Mar del Plata, Argentina, Comision Nacional De Investigaciones Espaciales, Argentina, 180 pp.
9. Muench, H. Stuart, "Large-Scale Disturbances in the Summertime Stratosphere," Journal of the Atmospheric Sciences, Vol. 25, No. 6, Nov. 1968, pp. 1108-1115.
10. Krumins, M. V., and Lyons, W. C., "Corrections for the Upper Atmosphere Temperatures Using a Thin Film Loop Mount," Technical Report 72-152, Naval Ordnance Laboratory, White Oak, Silver Spring, Md., June 1972, 38 pp.

11. Goddard Space Flight Center, "The Nimbus 4 User's Guide," Nimbus Project, U.S. National Aeronautical and Space Administration, Greenbelt, Md., 1970, 214 pp.
12. Goddard Space Flight Center, "The Nimbus 5 User's Guide," Nimbus Project, U.S. National Aeronautics and Space Administration, Greenbelt, Md., Nov. 1972, 162 pp.
13. McMillin, L. M., Wark, D. Q., Siomkajlo, J. M., Abel, P. G., Werbowetzki, A., Lauritson, L. A., Pritchard, J. A., Crossby, D. S., Woolf, H. M., Luebke, R. C., Weinreb, M. P., Fleming, H. E., Bittner, F. E., and Hayden, C. M., "Satellite Infrared Soundings From NOAA Spacecraft," NOAA Technical Report NESS 65, National Environmental Satellite Service, National Oceanic and Atmospheric Administration, U.S. Department of Commerce, Washington, D. C., Sept. 1973, 112 pp.
14. Quiroz, Roderick S., and Gelman, Melvyn E., "Direct Determination of the Thickness of Stratospheric Layers From Single-Channel Satellite Radiance Measurements," Monthly Weather Review, Vol. 100, No. 11, Nov. 1972, pp. 788-795.
15. NOAA, National Oceanographic and Atmospheric Administration, "Constant Pressure Charts 850-10 Mb," National Climatic Center MF-494-72,73, U.S. Department of Commerce, Asheville, N.C., 1972-1973.
16. Miller, Alvin J., "A Note on the Variability of Temperature as Indicated by Rocketsonde Thermistors," Journal of Applied Meteorology, Vol. 8, No. 1, Feb. 1969, pp. 172-174.
17. Miller, Alvin J., and Schmidlin, Francis J., "Rocketsonde Repeatability and Stratospheric Variability," Journal of Applied Meteorology, Vol. 10, No. 2, Apr. 1971, pp. 320-327.
18. Quiroz, Roderick S., (National Meteorological Center, National Weather Service, NOAA, Marlow Heights, Md.), "The Abnormal Stratosphere Studied With the Aid of Satellite Radiation Measurements," AIAA Paper No. 73-493, presented at the AIAA/AMS International Conference on the Impact of Aerospace Operations in the High Atmosphere, Denver, Colo., June 11-13, 1973, 8 pp.






Figure 1. Temperature measured at 55 km during 1972 at Ascension Island, Ft. Sherman, C.Z. and Thumba, India.




Figure 2. Relationships between satellite-measured radiances and thickness (a); and between radiance and temperature (b). The following relationships are shown: VTTPR channel 2 (NOAA 2) and SIRS channel 7 (Nimbus 4) for thickness between 100 to 5 mb; VTTPR channel 1 and SIRS channel 8 for thickness between 100 to 2 mb; SCR channel A (Nimbus 4) and SCR channel 2 (Nimbus 5) for thickness between 10 to 0.4 mb; SCR channel A and SCR channel 2 for temperature at 2 mb; and SCR channel A and SCR channel 2 for temperature at 5 mb. Radiance units  $10^{-7} \text{ J. cm}^{-2} \cdot \text{s}^{-1} \cdot (\text{ster})^{-1} \cdot (\text{cm}^{-1})^{-1}$ .




Figure 3a. Poker Flat, Alaska (65°07'N., 147°29'W.) analyzed values extracted from the 5-, 2-, and 0.4-mb charts for January to December 1972.




Figure 3b. Same as 3a for January to June 1973.




Figure 4a. Fort Churchill, Canada (58°44'N, 93°49'W.) analyzed values extracted from the 5-, 2-, and 0.4-mb charts for January to December 1972.




Figure 4b. Same as 4a for January to June 1973.






Figure 5a. Volgograd, U.S.S.R. (48°41'N, 44°21'E.) analyzed values extracted from the 5-, 2-, and 0.4-mb charts for January to December 1972.




Figure 5b. Same as 5a for January to June 1973.




Figure 6a. Wallops Island, Virginia (37°50'N, 75°29'W.) analyzed values extracted from the 5-, 2-, and 0.4-mb charts for January to December 1972.




Figure 6b. Same as 6a for January to June 1973.




Figure 7a. White Sands, N. Mexico ( $32^{\circ}23'N$ ,  $106^{\circ}29'W$ .) analyzed values extracted from the 5-, 2-, and 0.4-mb charts for January to December 1972.




Figure 7b. Same as 7a for January to June 1973.




Figure 8a. Antigua, W.I.A.S. (17°08'N, 61°47'W.) analyzed values extracted from the 5-, 2-, and 0.4-mb charts for January to December 1972.






Figure 8b. Same as 8a for January to June 1973.

# STATION MODEL AND REPORTING ROCKET STATIONS








Figure 9. Station Model and Reporting Rocket Stations.















































































































MAP 1000A  
 MAP 1000A AUG 71










UNIT: MILLIBARS  
 SCALE: 1:1000000  
 MAP 1000A AUG 71
























1000  
 1020  
 1040  
 1060  
 1080  
 1100  
 1120  
 1140  
 1160  
 1180  
 1200  
 1220  
 1240  
 1260  
 1280  
 1300  
 1320  
 1340  
 1360  
 1380  
 1400  
 1420  
 1440  
 1460  
 1480  
 1500  
 1520  
 1540  
 1560  
 1580  
 1600  
 1620  
 1640  
 1660  
 1680  
 1700  
 1720  
 1740  
 1760  
 1780  
 1800  
 1820  
 1840  
 1860  
 1880  
 1900  
 1920  
 1940  
 1960  
 1980  
 2000  
 2020  
 2040  
 2060  
 2080  
 2100  
 2120  
 2140  
 2160  
 2180  
 2200  
 2220  
 2240  
 2260  
 2280  
 2300  
 2320  
 2340  
 2360  
 2380  
 2400  
 2420  
 2440  
 2460  
 2480  
 2500  
 2520  
 2540  
 2560  
 2580  
 2600  
 2620  
 2640  
 2660  
 2680  
 2700  
 2720  
 2740  
 2760  
 2780  
 2800  
 2820  
 2840  
 2860  
 2880  
 2900  
 2920  
 2940  
 2960  
 2980  
 3000  
 3020  
 3040  
 3060  
 3080  
 3100  
 3120  
 3140  
 3160  
 3180  
 3200  
 3220  
 3240  
 3260  
 3280  
 3300  
 3320  
 3340  
 3360  
 3380  
 3400  
 3420  
 3440  
 3460  
 3480  
 3500  
 3520  
 3540  
 3560  
 3580  
 3600  
 3620  
 3640  
 3660  
 3680  
 3700  
 3720  
 3740  
 3760  
 3780  
 3800  
 3820  
 3840  
 3860  
 3880  
 3900  
 3920  
 3940  
 3960  
 3980  
 4000  
 4020  
 4040  
 4060  
 4080  
 4100  
 4120  
 4140  
 4160  
 4180  
 4200  
 4220  
 4240  
 4260  
 4280  
 4300  
 4320  
 4340  
 4360  
 4380  
 4400  
 4420  
 4440  
 4460  
 4480  
 4500  
 4520  
 4540  
 4560  
 4580  
 4600  
 4620  
 4640  
 4660  
 4680  
 4700  
 4720  
 4740  
 4760  
 4780  
 4800  
 4820  
 4840  
 4860  
 4880  
 4900  
 4920  
 4940  
 4960  
 4980  
 5000  
 5020  
 5040  
 5060  
 5080  
 5100  
 5120  
 5140  
 5160  
 5180  
 5200  
 5220  
 5240  
 5260  
 5280  
 5300  
 5320  
 5340  
 5360  
 5380  
 5400  
 5420  
 5440  
 5460  
 5480  
 5500  
 5520  
 5540  
 5560  
 5580  
 5600  
 5620  
 5640  
 5660  
 5680  
 5700  
 5720  
 5740  
 5760  
 5780  
 5800  
 5820  
 5840  
 5860  
 5880  
 5900  
 5920  
 5940  
 5960  
 5980  
 6000  
 6020  
 6040  
 6060  
 6080  
 6100  
 6120  
 6140  
 6160  
 6180  
 6200  
 6220  
 6240  
 6260  
 6280  
 6300  
 6320  
 6340  
 6360  
 6380  
 6400  
 6420  
 6440  
 6460  
 6480  
 6500  
 6520  
 6540  
 6560  
 6580  
 6600  
 6620  
 6640  
 6660  
 6680  
 6700  
 6720  
 6740  
 6760  
 6780  
 6800  
 6820  
 6840  
 6860  
 6880  
 6900  
 6920  
 6940  
 6960  
 6980  
 7000  
 7020  
 7040  
 7060  
 7080  
 7100  
 7120  
 7140  
 7160  
 7180  
 7200  
 7220  
 7240  
 7260  
 7280  
 7300  
 7320  
 7340  
 7360  
 7380  
 7400  
 7420  
 7440  
 7460  
 7480  
 7500  
 7520  
 7540  
 7560  
 7580  
 7600  
 7620  
 7640  
 7660  
 7680  
 7700  
 7720  
 7740  
 7760  
 7780  
 7800  
 7820  
 7840  
 7860  
 7880  
 7900  
 7920  
 7940  
 7960  
 7980  
 8000  
 8020  
 8040  
 8060  
 8080  
 8100  
 8120  
 8140  
 8160  
 8180  
 8200  
 8220  
 8240  
 8260  
 8280  
 8300  
 8320  
 8340  
 8360  
 8380  
 8400  
 8420  
 8440  
 8460  
 8480  
 8500  
 8520  
 8540  
 8560  
 8580  
 8600  
 8620  
 8640  
 8660  
 8680  
 8700  
 8720  
 8740  
 8760  
 8780  
 8800  
 8820  
 8840  
 8860  
 8880  
 8900  
 8920  
 8940  
 8960  
 8980  
 9000  
 9020  
 9040  
 9060  
 9080  
 9100  
 9120  
 9140  
 9160  
 9180  
 9200  
 9220  
 9240  
 9260  
 9280  
 9300  
 9320  
 9340  
 9360  
 9380  
 9400  
 9420  
 9440  
 9460  
 9480  
 9500  
 9520  
 9540  
 9560  
 9580  
 9600  
 9620  
 9640  
 9660  
 9680  
 9700  
 9720  
 9740  
 9760  
 9780  
 9800  
 9820  
 9840  
 9860  
 9880  
 9900  
 9920  
 9940  
 9960  
 9980  
 10000















- ▲ 12
- ▲ 13
- 14
- 15
- 16
- ⊗ 17












△ 10  
 ▲ 11  
 ● 12  
 ■ 13  
 □ 14

















10° 15° 20° 25° 30° 35° 40°  
 10° 15° 20° 25° 30° 35° 40°  
 10° 15° 20° 25° 30° 35° 40°  
 10° 15° 20° 25° 30° 35° 40°



SCALE 1:1,000,000  
POLAR STEREOGRAPHIC PROJECTION  
MAP 1000A AUG 71









MAP 1000A  
POLAR STEREOGRAPHIC PROJECTION  
MAP (GCS) JULY 71









5000 / 10000  
 POLAR STEREOGRAPHIC PROJECTION TIME 14 LUTON 80  
 MAP 1000A AUG 71









5 MB  
SEP 20, 1972

MAP 1000A  
MAP 1000A AND 71


- △ 18
- ▲ 19
- 20
- 21
- 22
- ⊠ 23
































NOAA J. HARRIS  
 POLAR STEREOGRAPHIC PROJECTION TIME OF LATITUDE 80°  
 MAP 1982A, MAY 71






















SCALE 1:100,000  
POLAR STEREOGRAPHIC PROJECTION TIME AT LUTTEREN HP  
MAP 1806A JULY 71
























NOV 22, 1972  
POLAR STATIONING INDICATIONS ARE AT 1000 HRS  
MAP 1000A AND 71











































18  
 19  
 20  
 21  
 22














MAP 1000A  
POLAR STEREOGRAPHIC PROJECTION TRUE AT LATITUDE 0°  
MAP 1000A, AUG 71




NOAA SYMBOLS  
 POLAR STEREOPROJECTION PROJECTIONS MADE AT LITTLETON, CO  
 MAP 1828A AUG 71










NOAA JPL/ARL  
POLAR STEREOGRAPHIC PROJECTION MAP OF LATITUDE 80°  
MAP 1828A AUG 71













UNIT: 1 MILLIBAR  
PACIFIC STANDARD TIME  
MAP 1800A JAN 71




1:1000000  
 POLAR STEREOPROJECTION  
 MAP 1808A AUG 71


UNIT: MILLIBARS  
 PRESSURE MEASUREMENTS: MEAN OF LAST FOUR  
 MAP: 1983A, AUG 71




2 MB  
 JAN 24, 1973

- △ 22
- △ 23
- 24
- 25
- 26
- ⊗ 27






DATE: 1 JAN 1973  
 TIME: 0000Z  
 MAP: 10000  
 MAP: 10000  
 MAP: 10000













UNIT: MILLIBARS  
PRESSURE: INTERPOLATED  
MAP: 1800A, 1800B, 1800C, 1800D, 1800E, 1800F, 1800G, 1800H, 1800I, 1800J, 1800K, 1800L, 1800M, 1800N, 1800O, 1800P, 1800Q, 1800R, 1800S, 1800T, 1800U, 1800V, 1800W, 1800X, 1800Y, 1800Z






































































UNIT: MILLIBARS  
 PRESSURE: 1000.0  
 MAP: 1000.0  
 MAP: 1000.0












190  
 191  
 192  
 193  
 194  
 195  
 196  
 197  
 198  
 199  
 200  
 201  
 202  
 203  
 204  
 205  
 206  
 207  
 208  
 209  
 210  
 211  
 212  
 213  
 214  
 215  
 216  
 217  
 218  
 219  
 220  
 221  
 222  
 223  
 224  
 225  
 226  
 227  
 228  
 229  
 230  
 231  
 232  
 233  
 234  
 235  
 236  
 237  
 238  
 239  
 240  
 241  
 242  
 243  
 244  
 245  
 246  
 247  
 248  
 249  
 250  
 251  
 252  
 253  
 254  
 255  
 256  
 257  
 258  
 259  
 260  
 261  
 262  
 263  
 264  
 265  
 266  
 267  
 268  
 269  
 270  
 271  
 272  
 273  
 274  
 275  
 276  
 277  
 278  
 279  
 280  
 281  
 282  
 283  
 284  
 285  
 286  
 287  
 288  
 289  
 290  
 291  
 292  
 293  
 294  
 295  
 296  
 297  
 298  
 299  
 300  
 301  
 302  
 303  
 304  
 305  
 306  
 307  
 308  
 309  
 310  
 311  
 312  
 313  
 314  
 315  
 316  
 317  
 318  
 319  
 320  
 321  
 322  
 323  
 324  
 325  
 326  
 327  
 328  
 329  
 330  
 331  
 332  
 333  
 334  
 335  
 336  
 337  
 338  
 339  
 340  
 341  
 342  
 343  
 344  
 345  
 346  
 347  
 348  
 349  
 350  
 351  
 352  
 353  
 354  
 355  
 356  
 357  
 358  
 359  
 360  
 361  
 362  
 363  
 364  
 365  
 366  
 367  
 368  
 369  
 370  
 371  
 372  
 373  
 374  
 375  
 376  
 377  
 378  
 379  
 380  
 381  
 382  
 383  
 384  
 385  
 386  
 387  
 388  
 389  
 390  
 391  
 392  
 393  
 394  
 395  
 396  
 397  
 398  
 399  
 400  
 401  
 402  
 403  
 404  
 405  
 406  
 407  
 4







MAP 1000A  
 POLAR STEREOGRAPHIC PROJECTION  
 DATE 1000A JUL 71


















UNIT 1: 1016.0  
UNIT 2: 1012.0  
UNIT 3: 1008.0  
UNIT 4: 1004.0  
UNIT 5: 1000.0  
UNIT 6: 996.0  
UNIT 7: 992.0  
UNIT 8: 988.0  
UNIT 9: 984.0  
UNIT 10: 980.0  
UNIT 11: 976.0  
UNIT 12: 972.0  
UNIT 13: 968.0  
UNIT 14: 964.0  
UNIT 15: 960.0  
UNIT 16: 956.0  
UNIT 17: 952.0  
UNIT 18: 948.0  
UNIT 19: 944.0  
UNIT 20: 940.0  
UNIT 21: 936.0  
UNIT 22: 932.0  
UNIT 23: 928.0  
UNIT 24: 924.0  
UNIT 25: 920.0  
UNIT 26: 916.0  
UNIT 27: 912.0  
UNIT 28: 908.0  
UNIT 29: 904.0  
UNIT 30: 900.0  
UNIT 31: 896.0  
UNIT 32: 892.0  
UNIT 33: 888.0  
UNIT 34: 884.0  
UNIT 35: 880.0  
UNIT 36: 876.0  
UNIT 37: 872.0  
UNIT 38: 868.0  
UNIT 39: 864.0  
UNIT 40: 860.0  
UNIT 41: 856.0  
UNIT 42: 852.0  
UNIT 43: 848.0  
UNIT 44: 844.0  
UNIT 45: 840.0  
UNIT 46: 836.0  
UNIT 47: 832.0  
UNIT 48: 828.0  
UNIT 49: 824.0  
UNIT 50: 820.0  
UNIT 51: 816.0  
UNIT 52: 812.0  
UNIT 53: 808.0  
UNIT 54: 804.0  
UNIT 55: 800.0  
UNIT 56: 796.0  
UNIT 57: 792.0  
UNIT 58: 788.0  
UNIT 59: 784.0  
UNIT 60: 780.0  
UNIT 61: 776.0  
UNIT 62: 772.0  
UNIT 63: 768.0  
UNIT 64: 764.0  
UNIT 65: 760.0  
UNIT 66: 756.0  
UNIT 67: 752.0  
UNIT 68: 748.0  
UNIT 69: 744.0  
UNIT 70: 740.0  
UNIT 71: 736.0  
UNIT 72: 732.0  
UNIT 73: 728.0  
UNIT 74: 724.0  
UNIT 75: 720.0  
UNIT 76: 716.0  
UNIT 77: 712.0  
UNIT 78: 708.0  
UNIT 79: 704.0  
UNIT 80: 700.0  
UNIT 81: 696.0  
UNIT 82: 692.0  
UNIT 83: 688.0  
UNIT 84: 684.0  
UNIT 85: 680.0  
UNIT 86: 676.0  
UNIT 87: 672.0  
UNIT 88: 668.0  
UNIT 89: 664.0  
UNIT 90: 660.0  
UNIT 91: 656.0  
UNIT 92: 652.0  
UNIT 93: 648.0  
UNIT 94: 644.0  
UNIT 95: 640.0  
UNIT 96: 636.0  
UNIT 97: 632.0  
UNIT 98: 628.0  
UNIT 99: 624.0  
UNIT 100: 620.0  
UNIT 101: 616.0  
UNIT 102: 612.0  
UNIT 103: 608.0  
UNIT 104: 604.0  
UNIT 105: 600.0  
UNIT 106: 596.0  
UNIT 107: 592.0  
UNIT 108: 588.0  
UNIT 109: 584.0  
UNIT 110: 580.0  
UNIT 111: 576.0  
UNIT 112: 572.0  
UNIT 113: 568.0  
UNIT 114: 564.0  
UNIT 115: 560.0  
UNIT 116: 556.0  
UNIT 117: 552.0  
UNIT 118: 548.0  
UNIT 119: 544.0  
UNIT 120: 540.0  
UNIT 121: 536.0  
UNIT 122: 532.0  
UNIT 123: 528.0  
UNIT 124: 524.0  
UNIT 125: 520.0  
UNIT 126: 516.0  
UNIT 127: 512.0  
UNIT 128: 508.0  
UNIT 129: 504.0  
UNIT 130: 500.0  
UNIT 131: 496.0  
UNIT 132: 492.0  
UNIT 133: 488.0  
UNIT 134: 484.0  
UNIT 135: 480.0  
UNIT 136: 476.0  
UNIT 137: 472.0  
UNIT 138: 468.0  
UNIT 139: 464.0  
UNIT 140: 460.0  
UNIT 141: 456.0  
UNIT 142: 452.0  
UNIT 143: 448.0  
UNIT 144: 444.0  
UNIT 145: 440.0  
UNIT 146: 436.0  
UNIT 147: 432.0  
UNIT 148: 428.0  
UNIT 149: 424.0  
UNIT 150: 420.0  
UNIT 151: 416.0  
UNIT 152: 412.0  
UNIT 153: 408.0  
UNIT 154: 404.0  
UNIT 155: 400.0  
UNIT 156: 396.0  
UNIT 157: 392.0  
UNIT 158: 388.0  
UNIT 159: 384.0  
UNIT 160: 380.0  
UNIT 161: 376.0  
UNIT 162: 372.0  
UNIT 163: 368.0  
UNIT 164: 364.0  
UNIT 165: 360.0  
UNIT 166: 356.0  
UNIT 167: 352.0  
UNIT 168: 348.0  
UNIT 169: 344.0  
UNIT 170: 340.0  
UNIT 171: 336.0  
UNIT 172: 332.0  
UNIT 173: 328.0  
UNIT 174: 324.0  
UNIT 175: 320.0  
UNIT 176: 316.0  
UNIT 177: 312.0  
UNIT 178: 308.0  
UNIT 179: 304.0  
UNIT 180: 300.0  
UNIT 181: 296.0  
UNIT 182: 292.0  
UNIT 183: 288.0  
UNIT 184: 284.0  
UNIT 185: 280.0  
UNIT 186: 276.0  
UNIT 187: 272.0  
UNIT 188: 268.0  
UNIT 189: 264.0  
UNIT 190: 260.0  
UNIT 191: 256.0  
UNIT 192: 252.0  
UNIT 193: 248.0  
UNIT 194: 244.0  
UNIT 195: 240.0  
UNIT 196: 236.0  
UNIT 197: 232.0  
UNIT 198: 228.0  
UNIT 199: 224.0  
UNIT 200: 220.0  
UNIT 201: 216.0  
UNIT 202: 212.0  
UNIT 203: 208.0  
UNIT 204: 204.0  
UNIT 205: 200.0  
UNIT 206: 196.0  
UNIT 207: 192.0  
UNIT 208: 188.0  
UNIT 209: 184.0  
UNIT 210: 180.0  
UNIT 211: 176.0  
UNIT 212: 172.0  
UNIT 213: 168.0  
UNIT 214: 164.0  
UNIT 215: 160.0  
UNIT 216: 156.0  
UNIT 217: 152.0  
UNIT 218: 148.0  
UNIT 219: 144.0  
UNIT 220: 140.0  
UNIT 221: 136.0  
UNIT 222: 132.0  
UNIT 223: 128.0  
UNIT 224: 124.0  
UNIT 225: 120.0  
UNIT 226: 116.0  
UNIT 227: 112.0  
UNIT 228: 108.0  
UNIT 229: 104.0  
UNIT 230: 100.0  
UNIT 231: 96.0  
UNIT 232: 92.0  
UNIT 233: 88.0  
UNIT 234: 84.0  
UNIT 235: 80.0  
UNIT 236: 76.0  
UNIT 237: 72.0  
UNIT 238: 68.0  
UNIT 239: 64.0  
UNIT 240: 60.0  
UNIT 241: 56.0  
UNIT 242: 52.0  
UNIT 243: 48.0  
UNIT 244: 44.0  
UNIT 245: 40.0  
UNIT 246: 36.0  
UNIT 247: 32.0  
UNIT 248: 28.0  
UNIT 249: 24.0  
UNIT 250: 20.0  
UNIT 251: 16.0  
UNIT 252: 12.0  
UNIT 253: 8.0  
UNIT 254: 4.0  
UNIT 255: 0.0













1000  
 950  
 900  
 850  
 800  
 750  
 700  
 650  
 600  
 550  
 500  
 450  
 400  
 350  
 300  
 250  
 200  
 150  
 100  
 50  
 0  
 -50  
 -100  
 -150  
 -200  
 -250  
 -300  
 -350  
 -400  
 -450  
 -500  
 -550  
 -600  
 -650  
 -700  
 -750  
 -800  
 -850  
 -900  
 -950  
 -1000  
 -1050  
 -1100  
 -1150  
 -1200  
 -1250  
 -1300  
 -1350  
 -1400  
 -1450  
 -1500  
 -1550  
 -1600  
 -1650  
 -1700  
 -1750  
 -1800  
 -1850  
 -1900  
 -1950  
 -2000  
 -2050  
 -2100  
 -2150  
 -2200  
 -2250  
 -2300  
 -2350  
 -2400  
 -2450  
 -2500  
 -2550  
 -2600  
 -2650  
 -2700  
 -2750  
 -2800  
 -2850  
 -2900  
 -2950  
 -3000  
 -3050  
 -3100  
 -3150  
 -3200  
 -3250  
 -3300  
 -3350  
 -3400  
 -3450  
 -3500  
 -3550  
 -3600  
 -3650  
 -3700  
 -3750  
 -3800  
 -3850  
 -3900  
 -3950  
 -4000  
 -4050  
 -4100  
 -4150  
 -4200  
 -4250  
 -4300  
 -4350  
 -4400  
 -4450  
 -4500  
 -4550  
 -4600  
 -4650  
 -4700  
 -4750  
 -4800  
 -4850  
 -4900  
 -4950  
 -5000  
 -5050  
 -5100  
 -5150  
 -5200  
 -5250  
 -5300  
 -5350  
 -5400  
 -5450  
 -5500  
 -5550  
 -5600  
 -5650  
 -5700  
 -5750  
 -5800  
 -5850  
 -5900  
 -5950  
 -6000  
 -6050  
 -6100  
 -6150  
 -6200  
 -6250  
 -6300  
 -6350  
 -6400  
 -6450  
 -6500  
 -6550  
 -6600  
 -6650  
 -6700  
 -6750  
 -6800  
 -6850  
 -6900  
 -6950  
 -7000  
 -7050  
 -7100  
 -7150  
 -7200  
 -7250  
 -7300  
 -7350  
 -7400  
 -7450  
 -7500  
 -7550  
 -7600  
 -7650  
 -7700  
 -7750  
 -7800  
 -7850  
 -7900  
 -7950  
 -8000  
 -8050  
 -8100  
 -8150  
 -8200  
 -8250  
 -8300  
 -8350  
 -8400  
 -8450  
 -8500  
 -8550  
 -8600  
 -8650  
 -8700  
 -8750  
 -8800  
 -8850  
 -8900  
 -8950  
 -9000  
 -9050  
 -9100  
 -9150  
 -9200  
 -9250  
 -9300  
 -9350  
 -9400  
 -9450  
 -9500  
 -9550  
 -9600  
 -9650  
 -9700  
 -9750  
 -9800  
 -9850  
 -9900  
 -9950  
 -10000





NOAA SYMBOLS  
 FOR WEATHER MAPS  
 MAP 1000A, AUG 71









POSTMASTER: If Undeliverable (Section 158  
Postal Manual) Do Not Return

*"The aeronautical and space activities of the United States shall be conducted so as to contribute . . . to the expansion of human knowledge of phenomena in the atmosphere and space. The Administration shall provide for the widest practicable and appropriate dissemination of information concerning its activities and the results thereof."*

—NATIONAL AERONAUTICS AND SPACE ACT OF 1958

## NASA SCIENTIFIC AND TECHNICAL PUBLICATIONS

**TECHNICAL REPORTS:** Scientific and technical information considered important, complete, and a lasting contribution to existing knowledge.

**TECHNICAL NOTES:** Information less broad in scope but nevertheless of importance as a contribution to existing knowledge.

**TECHNICAL MEMORANDUMS:** Information receiving limited distribution because of preliminary data, security classification, or other reasons. Also includes conference proceedings with either limited or unlimited distribution.

**CONTRACTOR REPORTS:** Scientific and technical information generated under a NASA contract or grant and considered an important contribution to existing knowledge.

**TECHNICAL TRANSLATIONS:** Information published in a foreign language considered to merit NASA distribution in English.

**SPECIAL PUBLICATIONS:** Information derived from or of value to NASA activities. Publications include final reports of major projects, monographs, data compilations, handbooks, sourcebooks, and special bibliographies.

**TECHNOLOGY UTILIZATION PUBLICATIONS:** Information on technology used by NASA that may be of particular interest in commercial and other non-aerospace applications. Publications include Tech Briefs, Technology Utilization Reports and Technology Surveys.

Details on the availability of these publications may be obtained from:

SCIENTIFIC AND TECHNICAL INFORMATION OFFICE

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

Washington, D.C. 20546